

A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors

Hildur V. Colot^{*†}, Gyungsoon Park^{†‡}, Gloria E. Turner[§], Carol Ringelberg^{*}, Christopher M. Crew^{*†¶}, Liubov Litvinkova[‡], Richard L. Weiss[§], Katherine A. Borkovich[‡], and Jay C. Dunlap^{*||}

^{*}Department of Genetics, Dartmouth Medical School, HB7400, Hanover, NH 03755; [†]Department of Plant Pathology, University of California, Riverside, CA 92521; and [§]Department of Chemistry and Biochemistry, 405 Hilgard Avenue, University of California, Los Angeles, CA 90095

Edited by David D. Perkins, Stanford University, Stanford, CA, and approved March 29, 2006 (received for review February 21, 2006)

The low rate of homologous recombination exhibited by wild-type strains of filamentous fungi has hindered development of high-throughput gene knockout procedures for this group of organisms. In this study, we describe a method for rapidly creating knockout mutants in which we make use of yeast recombinational cloning, *Neurospora* mutant strains deficient in nonhomologous end-joining DNA repair, custom-written software tools, and robotics. To illustrate our approach, we have created strains bearing deletions of 103 *Neurospora* genes encoding transcription factors. Characterization of strains during growth and both asexual and sexual development revealed phenotypes for 43% of the deletion mutants, with more than half of these strains possessing multiple defects. Overall, the methodology, which achieves high-throughput gene disruption at an efficiency >90% in this filamentous fungus, promises to be applicable to other eukaryotic organisms that have a low frequency of homologous recombination.

filamentous fungi | functional genomics | knockouts | recombinational cloning

Historically, it has proven difficult to perform gene knockouts on a large scale in filamentous fungi. Unlike *Saccharomyces cerevisiae*, wild-type *Neurospora* strains exhibit a low frequency of homologous recombination after transformation, typically <10%, even when deletion cassettes contain long stretches of flanking sequence. The predominance of ectopic insertions dictates that a large number of transformants be screened. Of the many published procedures for creating knockout constructs (e.g., refs. 1–6), none lends itself well to a reliable high-throughput approach. To fill this need, we have developed a scheme that takes advantage of recombinational cloning in yeast (7, 8), a method that bypasses traditional restriction digestion and ligation. The components of the final construct are synthesized individually with short overlapping ends by PCR and cotransformed into yeast for assembly by the recombination machinery. An additional significant advance is the creation of *Neurospora* recipient strains in which ectopic insertions are virtually eliminated; Inoue and coworkers (9) recently demonstrated that mutation of either of two *Neurospora* genes required for nonhomologous end-joining DNA repair (*mus-51* and *mus-52*) results in a dramatic increase in the frequency of homologous recombination. We have combined our method for cassette synthesis, the use of Δ *mus-51* and Δ *mus-52* strains, and additional procedures including custom-designed software tools, into a successful high-throughput gene deletion strategy for filamentous fungi. Here we report application of this technology to mutation of 103 genes encoding transcription factors in *Neurospora*.

The filamentous fungus *Neurospora crassa* is a model for many fungal species that are important animal and plant pathogens. *Neurospora* possesses a more complex life cycle than yeasts, with three different sporulation pathways and production of at least 28 different cell types (10, 11). Although *Neurospora* has been

studied for >60 years by classical and molecular genetics, relatively little is known about transcriptional regulation in this organism. The availability of the complete *Neurospora* genome sequence (12) has allowed annotation of 182 transcription factor genes (not including general factors that regulate RNA polymerase; see ref. 11). The majority of factors belong to the Zn(II)₂Cys₆ fungal binuclear cluster family, whereas others are members of the C2H2 zinc finger, GATA, bHLH, B-ZIP, CBF CAAT-binding factor, homeobox, RING finger, PHD finger, MIZ zinc finger, and other families (11). Only 21 transcription factor genes have been characterized in *Neurospora*, and fewer yet have been studied in any other filamentous fungal system. The genes that have been studied in *Neurospora* play roles in metabolism and during development, including nutrient utilization or uptake (*acu-15*, *cpc-1*, *nit-2*, *nit-4*, *pco-1*, *nuc-1*, *cys-3*, *qa-1F*, and *sre*; see refs. 13–21), vegetative growth and macroconidiation (*fl*, *rca-1*, and *rco-1*; see refs. 22 and 23), light responses and the circadian rhythm (*wc-1* and *wc-2*; see refs. 24 and 25), sexual development (*asd-4*, *asm-1*, and *pp-1*; see refs. 26–28), DNA repair (*uvs-2*; see ref. 29), and mating type functions (*mat A-1*, *mat A-3*, and *mat a-1*; see refs. 30 and 31). There are many other processes for which basic information regarding transcriptional control mechanisms is totally lacking. Therefore, a large-scale functional study of transcription factors in *Neurospora* is timely and will provide a foundation for understanding complex modes of gene regulation in filamentous fungi.

Results

The overall strategy for high-throughput generation of gene deletions required a series of methodological advances for the construction of the deletion cassette DNA. In addition, recipient strains were created for transformation (*mus-51::bar* and *mus-52::bar*; see *Materials and Methods*). Because the marker for our deletion cassettes is *hph* (see below), we were unable to use the *mus-51/52::hph* deletion strains reported by Ninomiya *et al.* (9).

Primer Design, Optimization, and Production of Deletion Cassettes.

The cassettes contain (in order, 5' to 3') 1–1.3 kb of 5' sequence flanking the ORF to be deleted (target gene ORF); the selectable marker *hph* [the gene for hygromycin B phosphotransferase (32), which confers hygromycin resistance] driven by the *trpC*

Conflict of interest statement: No conflicts declared.

This paper was submitted directly (Track II) to the PNAS office.

Freely available online through the PNAS open access option.

Abbreviation: VM, Vogel's minimal medium.

[†]H.V.C. and G.P. contributed equally to this work.

[¶]Present address: School of Psychology, Georgia Institute of Technology, 654 Cherry Street, Atlanta, GA 30332-0170.

^{||}To whom correspondence should be addressed. E-mail: jay.c.dunlap@dartmouth.edu.

© 2006 by The National Academy of Sciences of the USA




Fig. 1. Strategy for creating deletion constructs. 5' and 3' flank fragments are amplified separately from genomic DNA with primers 5f + 5r and 3f + 3r. Primers 5r and 3f incorporate MmeI sites (M) and have 5' tails homologous to the *hph* cassette, whereas those for 5f and 3r are homologous to the vector. The two flanks are cotransformed into yeast along with the *hph* cassette and gapped yeast shuttle vector. Homologous recombination creates the circular construct and the final linear deletion cassette is amplified from the pooled yeast DNA with primers 5f and 3r. *hph* is transcribed in the antisense direction relative to the target gene.

promoter (33) and flanked by engineered MmeI restriction sites; and 1–1.3 kb of 3' flanking sequence for the target gene ORF.

The method for construction of deletion cassettes is presented in Fig. 1. Briefly, four DNA fragments comprising the 5' and 3' DNA sequences flanking the target gene, the *hph* cassette, and a gapped yeast vector, respectively, were assembled in yeast using the endogenous homologous recombination system, and the final linear deletion cassette was amplified from the resulting construct by PCR (see Fig. 1 and *Materials and Methods*). To achieve this, four PCR primers, designated 5f, 5r, 3f, and 3r, were synthesized for each knockout cassette. The common 29-nt 5' portions of the primers are shown in *Supporting Text*, which is published as supporting information on the PNAS web site; to select the gene-specific 20-nt portions of the primers, we used a custom-built software application (see *Materials and Methods*). This program successfully designed primers for 10,047 (96.8%) of the 10,620 ORFs predicted by Assembly 7 of the *Neurospora* genome sequence (www.broad.mit.edu/annotation/fungi/neurospora_crassa_7/index.html). Primer design failed for 16 genes because of their proximity to the ends of contigs. Modest alterations in the selection parameters enabled successful primer design for most of the other failures (data not shown).

MmeI sites were designed into the junctions between the flanks and the *hph* cassette (Fig. 1) to allow the adjacent gene-specific 20-bp sequences to serve as molecular bar codes. Digestion of genomic DNA by MmeI (which cuts 20 base pairs downstream of its recognition sequence) followed by ligation-mediated PCR will create products that can be amplified and then either sequenced or used in standard competition experiments (34–36).

In the development phase of this work, deletion cassettes were made with the *hph* gene driven by the *Ashbya gossypii* TEF promoter (pTEF); pTEF drives expression in both yeast and *Neurospora*, thereby allowing direct selection of yeast cells

harboring the correctly assembled constructs on medium containing hygromycin B. Although sufficient for creation of deletion strains (e.g., C. A. Jones, S. E. Greer-Phillips, and K.A.B., unpublished work), the weak activity of pTEF in *Neurospora* led to slow growth and low numbers of transformants at the hygromycin concentration necessary for inhibiting growth of nontransformed cells. Therefore, to ensure dependable selection of true *Neurospora* transformants, we settled on the widely used and highly expressed *Aspergillus nidulans trpC* promoter-*hph* cassette (33) as our selectable marker for *Neurospora* deletion constructs. The yeast vector contains the *URA3* gene, and selection is done on medium lacking uracil. DNA made from pooled yeast transformants serves as the template for specific amplification of the correct deletion cassette.

Before Inoue and coworkers published their findings regarding the *mus-51* and *mus-52* deletion strains (9), we explored methods for increasing the frequency of homologous recombination among transformants. Deletion cassette constructs were generated by yeast recombinational cloning as shown in Fig. 1, except that the flank fragments were 3 kb long. Split-marker fragments (37) were then amplified by PCR from the pooled yeast DNA (see Fig. 4, which is published as supporting information on the PNAS web site) and cotransformed into *Neurospora*. Combining our method for cassette construction with the published split-marker technique gave results that were a significant improvement over existing approaches. On average, 44% of the transformants obtained with this strategy had the proper gene replacement and, as determined by Southern analysis, 68% of those were free of ectopic insertions (results for 16 individual genes vary widely and are shown in Table 2, which is published as supporting information on the PNAS web site). Our subsequent adoption of Δ *mus-51* and Δ *mus-52*, along with deletion cassettes with 1-kb flanks, rendered the split-marker approach unnecessary for the high-throughput deletion project. However, it remains a useful strategy for transforming strains not deficient in *mus-51* or *mus-52*, especially when combined with our method for creating deletion cassettes.

Subsequent to the creation of the transcription factor deletion cassettes, this method was used to produce deletion cassettes for >9,600 *Neurospora* genes. The overall success rate for synthesis of flank fragments is $\approx 97\%$, whereas that for PCR amplification of the final deletion cassettes from the mixed yeast DNA pools is 90–95%. Electroporation of *Neurospora* with the final cassettes used for disrupting the transcription factors worked well over at least a 5-fold range of input DNA amounts, typically 200–1,000 ng.

Generation of Transcription Factor Deletion Mutants. The 103 transcription factor genes chosen to illustrate the utility of this procedure were previously annotated (11) by BLAST searches (38) by using Assembly 3 of the *Neurospora* genome sequence (www.broad.mit.edu/annotation/fungi/neurospora_crassa_3/index.html). Genes with E values $< 10^{-5}$ were accepted as members of a particular class. For initial knockout experiments, genes representing all of the annotated groups (11) were chosen. Cassettes were made and used to transform Δ *mus* strains as described above and in *Materials and Methods*, and the heterokaryotic primary transformants bearing putative gene knockouts were selected on the basis of hygromycin resistance.

Southern blot analysis of 627 primary transformants (six to seven transformants per gene) showed that homologous recombination without any ectopic insertion occurred in 98% of the cases. Ectopic insertion of knockout cassette DNA in addition to homologous recombination was observed in 0.8% of transformants, but stable homokaryons free of ectopic sequences were eventually obtained from these transformants after crossing (Table 1). Ectopic insertions without concurrent homologous

Table 1. Results of Southern blot analysis of 627 primary transformants

	HR* with no ectopic†	HR with ectopic‡	Ectopic with no HR§	Unusual recombination¶
Number	613	5	6	3
Percent of total	98	0.8	0.8	0.4

*HR, Homologous recombination.

†Transformants with bands indicative of gene replacement as well as wild type, but with no additional bands indicative of ectopic insertion.

‡Transformants with bands indicative of gene replacement and wild type, as well as additional bands.

§Transformants with no bands indicative of gene replacement but with both wild type and additional bands.

¶Transformants with no bands indicative of gene replacement or wild type but only extra bands.

recombination, as well as abnormal recombination events, were observed in 1.2% of the transformants (Table 1).

Primary transformants were crossed to wild-type females to obtain homokaryotic knockout mutants free of the *mus* mutation. Although 99 of the transcription factor genes yielded homokaryotic deletion progeny, no hygromycin-resistant progeny were isolated for NCU00340 (*pp-1*), NCU01345 (*asl-1*), NCU01459 (*asl-2*), and NCU04050 (*cpc-1*).

Phenotypic Analysis of Transcription Factor Deletion Strains. A substantial fraction of the viable mutants displayed visible phenotypes; the results indicate that 40 of the 99 genes studied are required for normal vegetative or sexual growth and development in *Neurospora* (Fig. 2). Table 3, which is published as supporting information on the PNAS web site, provides a complete list of genes deleted and the results of the phenotypic analyses, as well as names applied to mutants exhibiting novel phenotypes. Further details and photographs of the mutant strains are available at www.broad.mit.edu/annotation/genome/neurospora/Alleles.html.

Basal hyphal extension and asexual development. *Neurospora* grows by apical extension, branching, and fusion of basal hyphae to form a colony. Nutrient deprivation or desiccation leads to differentiation of aerial hyphae from these basal hyphae. The ends of aerial hyphae form budding structures called conidiophores that give rise to mature vegetative spores, the macroconidia (39).

In our study, 20 mutants exhibited significantly reduced basal hyphal extension rates compared to wild type as measured in race tubes (see *Supporting Text*), whereas five mutants had faster growth rates (Fig. 2 and Table 3). Most mutants with defects in extension of basal hyphae also demonstrated abnormalities during macroconidiation (15 strains; Fig. 2). A positive correlation between production of normal levels of mature macroconidia and the height of aerial hyphae was observed for many mutants (data not shown). For example, the Δ *fluffy* (*fl*):NCU08726 mutant exhibited short aerial hyphae and a block in macroconidiation as previously reported (40, 41). However, nine mutant strains with short aerial hyphae formed macroconidia normally (Table 3). Five mutants extended longer aerial hyphae than wild type (Fig. 2). Of particular interest, two mutants (Δ *vad-6*:NCU09205 and Δ *bek-2*:NCU07139; see below) exhibited both faster basal hyphal extension rates and longer aerial hyphae than wild type (Fig. 2).

The Δ *kaleidoscope-1* (*kal-1*:NCU03593) mutant was so named because of its radial and variable colony appearance under different growth conditions. The Δ *kal-1* mutant has a slower basal hyphal extension rate than wild type at 30°C and 37°C, but not at 25°C (Fig. 3A and data not shown). When 2% yeast extract was added to the medium, extension of basal hyphae was decreased in wild type but increased in the Δ *kal-1* strain (Fig.




Fig. 2. Venn diagram showing the distribution of transcription factor knockout mutants with observed phenotypes. Mutants are represented by the NCU numbers of the deleted genes. Font color indicates gene family (see *Inset* table). Miscellaneous genes (pink) are from three classes: RING-type zinc finger (NCU06411), CBF CAAT-binding factor (NCU02017), and homeobox (NCU00097 and NCU03593). Genes showing ascospore lethality are shown in the oval. The numbers of transcription factor mutants analyzed and the numbers with phenotypes are listed in the *Inset* table. Knockout mutants indicated with * or + exhibited greater basal hyphal extension rates or aerial hyphae heights than wild type, respectively. For the wild-type controls, the basal hyphae extension rate was 65–80 mm per day, whereas the height of aerial hyphae achieved in 3 days was 30–45 mm. The three photographs show a perithecium (Upper), a conidiophore (Lower Right), and basal hyphae (Lower Left).

3A). Δ *kal-1* mutants also exhibit increased branching of basal hyphae and shorter aerial hyphae (Fig. 3A and Table 3).

Sexual development. *Neurospora* differentiates female reproductive structures (protoperithecia) in response to nitrogen starvation. Fertilization is accomplished by chemotropic growth of a specialized female hypha toward a male cell (typically a macroconidium) of opposite mating type, transport of the male nucleus into the protoperithecium, cell proliferation, and karyogamy. Meiosis ensues and the protoperithecium develops into the mature fruiting body (perithecium) containing ascospores (42).

All 99 viable knockout mutants exhibited normal male fertility (data not shown). In contrast, 14 of the transcription factors studied are required for various stages of female sexual development and ascospore production (Fig. 2 and Table 3). Nine of the mutants that do not produce perithecia also form few or no protoperithecia. The arrested development (Δ *adv-1*:NCU07392) mutant is blocked in protoperithecial formation (as well as basal hyphal extension and asexual development; see below), in contrast to strains lacking the homologous gene *pro1* in *Sordaria macrospora* (43). Submerged protoperithecia were observed in the Δ *sub1*:NCU01154 mutant. Several mutants develop aberrant perithecia, including defective beaks (Δ *bek-1*:NCU00097 and Δ *bek-2*; see above; Fig. 3B) and less melanization (Δ *mel-1*:NCU04561). Of these three mutants, only Δ *mel-1* is able to efficiently eject ascospores. Interestingly, the Δ *nit-4*:NCU08294




Fig. 3. Colony morphology and asexual and sexual development of transcription factor knockout mutants. (A) Colony morphology of wild type and $\Delta kal-1$:NCU03593. Strains were grown for 24 h on VM or VM + yeast extract at 25°C and 37°C. The colony edge images show basal vegetative hyphae at $\times 11$. (B) Transcription factor mutant exhibiting aberrant development of perithecia. Images at $\times 77$ were taken 7 days after fertilization of protoperithecia with opposite mating type conidia. The arrow indicates a beak in wild type; this structure does not form in the mutant.

mutant forms more protoperithecia and perithecia than wild type (data not shown).

Eight transcription factor knockout mutants possess significant defects in all three phenotypes: basal hyphal extension, asexual development, and sexual development (Fig. 2). In contrast, $\Delta bek-2$ has a defect in sexual development but demonstrates increased extension of both basal and aerial hyphae (Fig. 2). With the exception of *all development altered* (*ada-2*:NCU02017, *ada-4*:NCU03320, and *adv-1*:NCU07392, these genes and their closest homologues have no reported functions in any filamentous fungal system.

Discussion

Methodological advances reported here significantly facilitate the high-throughput production of deletion mutants. We found recombinational cloning in yeast, a well characterized and widely used technique, to be the most suitable method for high-throughput synthesis of deletion constructs. Most steps are performed on a pipetting robot, allowing the creation of 400–600 constructs per week. Cassettes are electroporated into the *mus* deletion mutants in a 96-well format. Strains and processes are tracked by using a custom-written LABORATORY INFORMATION MANAGEMENT SYSTEM (LIMS; see *Supporting Text*), and streamlined methods for preliminary phenotypic analysis have been developed. Both the general procedures developed for *Neurospora* (e.g., the use of *mus-51/52* deletions to reduce ectopic insertions and a strong promoter to drive selectable marker expression) and the technical developments (primer selection and Southern analysis programs; high-throughput cassette construction, preparation, and transformation) should be readily adaptable to other organisms. In addition, this powerful methodology can be used to generate strains with tagged or

mutant alleles or promoter replacements at the endogenous locus of any given gene.

The 99 viable transcription factor mutants were analyzed for a variety of characteristics during vegetative and sexual growth and development. A total of 40 mutants displayed at least one defining phenotype during our testing. Among the genes producing phenotypes, only four had been studied previously in *Neurospora* [*wc-2*:NCU00902, *acu-15*:NCU06656, *fluffy* (*fl*), and *nit-4*], and 14 others have characterized homologues in other fungi.

Our results revealed that highly conserved transcription factors can play different roles in various fungi. For example, *Sordaria macrospora pro1* mutants are able to form protoperithecia, but not perithecia (43). In contrast, the *Aspergillus nidulans* Pro1 homologue RosA is required for sexual development at low carbon levels (44). Our analysis of the *Neurospora* homologue of *pro1*, *adv-1*, demonstrates that it is required for normal protoperithecial differentiation at both high and low sucrose concentrations. Thus, *adv-1* is likely to be required during an earlier stage of female sexual development than *pro1*, and it does not possess a carbon concentration-dependent function, as observed for *rosA*. In addition, *adv-1* is also essential for normal vegetative growth and development in *Neurospora*.

That no viable ascospore progeny were observed for four transcription factor mutants ($\Delta pp-1$:NCU00340, $\Delta cpc-1$:NCU04050, $\Delta asl-1$:NCU01345, and $\Delta asl-2$:NCU01459) suggests these factors are required for ascospore germination and/or perform some other essential function during the life cycle. Similar results were obtained during analysis of *S. cerevisiae* transcription factor genes, with 3.4% found to be essential for viability in rich medium (45). Interestingly, with the exception of *pp-1*, all of the ascospore lethal *Neurospora* genes are members of the B-ZIP family (11). The results for *pp-1* are supported by a recent report that ascospores carrying a deletion of *pp-1* are inviable, but that $\Delta pp-1$ strains can be maintained in the vegetative phase (26). Viable $\Delta cpc-1$ mutants have been reported (46), and we were eventually able to isolate $\Delta cpc-1$ strains by using the microconidiation procedure in the vegetative phase to bypass the need for a sexual cross (data not shown; see ref. 47).

Our work uncovered an important regulator of asexual growth and development in *Neurospora*, *kaleidoscope-1* (*kal-1*), which encodes a homeobox-containing transcription factor. *kal-1* is homologous to *Podospira anserina pah1*, a gene required for normal microconidiation and hyphal branching (48). The $\Delta kal-1$ mutation is highly pleiotropic, leading to substantial changes in colony morphology and conidial development (Fig. 3A). However, supplementation with yeast extract increases apical extension and conidiation of the *kal-1* mutant, indicating a possible role for *kal-1* in nutrient metabolism or sensing. Despite their extensive vegetative defects, *kal-1* mutants are female-fertile. Two additional homeobox transcription factor genes, *bek-1* and NCU03070, were also examined in our study; *bek-1* was found to be required for normal development of perithecial beaks during sexual development. Homeobox transcription factors are well known as regulators of cell proliferation, differentiation, and pattern formation in mammals (49–51); our analysis supports conservation of developmental roles for homeobox factors in *Neurospora*.

It is notable that 59 of the transcription factor genes studied here do not possess obvious functions during growth and development, suggesting they may have roles under environmental conditions not included in our analyses. Some of the genes may also possess overlapping functions. Functional redundancy is observed for many proteins in eukaryotes, including transcription factors (52–54), and can be preserved over long periods of evolution because the compensatory activities of related genes can buffer the effects of deleterious mutations (55). Remarkably,

numerous transcription factor genes in our study (33%) are similar to another *Neurospora* transcription factor (e^{-64} to e^{-5}), and the majority of these proteins belong to the Zn(II)₂Cys₆ fungal binuclear cluster family (11). In addition, many (79%) did not yield obvious phenotypes when mutated. Mutation of multiple homologous transcription factor genes will be necessary to comprehend complex gene regulation in *Neurospora*.

Characterization of the null phenotype is a requisite step in determining the functions of a gene, and the accessibility of a complete collection of mutants would begin to unlock the secrets of any genome. We have reported here the development and implementation of a scheme for high-throughput production of targeted gene deletions in *Neurospora* that should be widely applicable to the filamentous fungi and beyond. We have used these methods to systematically delete genes encoding 103 transcription factors, and the resulting strains have provided a window into novel processes controlling vegetative growth, as well as sexual and asexual development.

Materials and Methods

Neurospora Strains. Strains FGSC 4200 (*mat a*), FGSC 2489 (*mat A*), FGSC 6103 (*his-3 mat A*), FGSC 4317 (*fl mat A*), and FGSC 4347 (*fl mat a*) were obtained from the Fungal Genetics Stock Center, Kansas City, KS. Strains FGSC 9717 (*mus-51::bar his-3 mat A*), FGSC 9718 (*mus-51::bar mat a*), FGSC 9719 (*mus-52::bar mat a*), and FGSC 9720 (*mus-52::bar his-3 mat A*) were created in this study.

Primer Design and Synthesis of Deletion Cassettes. A software application written by John Jones (John Jones Consulting, Moreno Valley, CA) was designed to retrieve regions adjacent to each ORF and pass them to PRIMER3 (http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi), which would automatically select a list of candidate primers with designated parameters (see *Supporting Text*). For gene-specific flanks, the program searched a 1,500-bp region on either side of the ORF for primers that would synthesize a 1- to 1.3-kb fragment, thus allowing a gap between the ORF and either flank fragment of 0–500 bp. The resulting suggested primers were then tested for the absence of MmeI restriction sites (see above), uniqueness in the genome [by BLAST; (38)], and suitable GC content (to minimize primer-dimer formation), and the top-rated primer pair was selected. Primers were supplied in 96-well plates (Illumina, San Diego) and were diluted and mixed in three pairs for each gene by a Biomek NX robot (Beckman): 5f + 5r and 3f + 3r for PCR synthesis of the 5' and 3' flank fragments, respectively, and 5f + 3r for production of the final cassettes. The flanks were synthesized from genomic *Neurospora* DNA using LA Taq (TaKaRa; exact details of all procedures are in *Supporting Text*).

Yeast Transformation. For the 103 transcription factor gene knockouts, yeast strain FY834 (56) was transformed with both flank fragments, the *hph* cassette, and gapped plasmid [pRS426 (57) digested with XbaI and EcoRI] in 96-well plates on the robot. A lithium chloride/polyethylene glycol procedure developed for a 96-well format was used essentially as described (58), except that the selection for transformants was done in liquid medium lacking uracil.

Preparation of Yeast DNA and Generation of Deletion Cassettes by PCR. Yeast DNA was prepared from the liquid cultures on the robot with the Puregene Yeast DNA kit (Gentra Systems) with slight modifications: Zymolyase T-100 (Seikagaku America, Falmouth, MA) was used and, in place of the final precipitation step, CleanSEQ magnetic beads (Agencourt, Beverly, MA) were used to bind the DNA. For the synthesis of full-length cassettes, yeast DNA and the primer pair mixture 5f + 3r for each gene were added by the robot to a PCR mix containing LA Taq.

Construction of the *mus-51* and *mus-52* Deletion Mutant Strains. Yeast recombinational cloning was performed to construct deletion cassettes with the *bar* cassette [which confers resistance to phosphinothricin (59, 60)] flanked by 3 kb upstream and 3 kb downstream of the ORF of either *mus-51* or *mus-52*. The 7-kb deletion cassette fragment was generated by using the appropriate 5f and 3r primers with the yeast DNA as a template for PCR. The cassettes were transformed by electroporation into wild-type *Neurospora* strain FGSC 4200, as described (61, 62). Selection of transformants, Southern analysis, and sexual crosses to obtain homokaryons are described in *Supporting Text*. Four different genotypes of Δ *mus-51* and Δ *mus-52* were obtained (see above); all four showed normal vegetative growth and sexual fertility, consistent with the previous study (9).

High-Throughput Generation of 103 Transcription Factor Deletion Mutants. Conidia from either Δ *mus-51* (FGSC 9718) or Δ *mus-52* (FGSC 9719) were transformed by electroporation with each knockout cassette, allowed to recover in Vogel's minimal medium (VM; ref. 63) supplemented with yeast extract and histidine and plated in regeneration agar on sorbose plates (61) containing hygromycin B, yeast extract, and histidine. After incubation for 5 days at 30°C in the dark, colonies were picked onto VM slants containing hygromycin B.

Six or more transformants for each gene were subjected to Southern blot analysis, as described (64), with some modifications. A program was developed to allow automated identification of appropriate restriction enzymes to use for Southern blot analysis of gene replacement mutants (developed by John Jones; details in *Supporting Text*). The DIG High Prime labeling kit (Roche Applied Science, Indianapolis) was used for labeling full-length knockout cassette fragment probes.

Heterokaryotic transformants confirmed by Southern analysis to contain both wild-type and deletion mutant nuclei were crossed to wild-type (FGSC 2489) or *his-3* (FGSC 6103) females. Ascospore progeny resistant to hygromycin were selected and tested for the *mus* deletion mutation by assessing sensitivity to phosphinothricin. Confirmation of homokaryons by Southern blot analysis, and determination of mating type and *his-3* alleles were as described in *Supporting Text*.

Phenotypic Analysis. Three major phenotypes were analyzed in verified transcription factor knockout mutants in an otherwise wild-type background: linear growth rates of basal hyphae, asexual development, and sexual development. Detailed procedures are presented in *Supporting Text*. Linear growth rates for basal hyphae were measured by using VM agar race tubes. Colony morphology and conidiation were assessed after inoculation of VM and VM + yeast extract plates and incubation at 25°C or 37°C. VM slant tubes inoculated with strains, incubated at 25°C for 3 days and then at room temperature for 3–5 days, were used to analyze aerial hyphae production, conidiation, and pigmentation. Conidiation was also assessed on plates. Extension of aerial hyphae in standing liquid cultures was measured after incubation for 72 h at 25°C.

Sexual development analysis was performed after inoculation of strains on synthetic crossing medium plates containing 0.1% or 1.5% sucrose and incubation at room temperature. Production of protoperithecia was scored after 7–8 days. At this time, plates were fertilized with opposite mating type wild-type strains and perithecial formation determined after an additional 7–8 days. Ascospore ejection was assessed 2 weeks after fertilization.

We thank John Jones for software design and LIMS implementation; Svetlana Krystofova and Suzanne Phillips for help with *mus* strain construction; Victoria Copeland for excellent technical assistance; and the following students who participated in the *Neurospora* Genetics and Genomics Summer Research Institute (NGGSRI) at the University of

California, Los Angeles, for phenotypic analysis: Cynthia Aguirre, Eliana Alcantar, Andrea Cahuantzi, Natalie Cornejo, Zachary W. Cue, Evelyn De Los Santos, Anthony Dualo, Thomas J. Duneheiw, Mina El-Masry, Jonathan Finley, Lizette C. Flores, Christopher Fonseca, Rukhsana A. Khan, Carolyn Kingsley, Juan Lupercio, Criseyda Mar-

tinez, Rosaura Ochoa, Olufisayo Oke, Cam M. Phu, Chloe Rivera, Michael Smith, Desiree L. Tejada, Tuan D. Tran, and Jackelyn Valladares. This work was supported by National Institutes of Health Grant P01 GM068087. The NIGMS program was supported by National Institutes of Health Grants 5 R25 GM050067 and 5 R25 GM055052.

- Wendland, J. (2003) *Curr. Genet.* **44**, 115–123.
- Yang, L., Ukil, L., Osmani, A., Nahm, F., Davies, J., De Souza, C. P., Dou, X., Perez-Balaguer, A. & Osmani, S. A. (2004) *Eukaryot. Cell* **3**, 1359–1362.
- Chaverroche, M. K., Ghigo, J. M. & d'Enfert, C. (2000) *Nucleic Acids Res.* **28**, E97.
- Yu, J. H., Hamari, Z., Han, K. H., Seo, J. A., Reyes-Dominguez, Y. & Scazzocchio, C. (2004) *Fungal Genet. Biol.* **41**, 973–981.
- Kuwayama, H., Obara, S., Morio, T., Katoh, M., Urushihara, H. & Tanaka, Y. (2002) *Nucleic Acids Res.* **30**, E2.
- Aronson, B. D., Lindgren, K. M., Dunlap, J. C. & Loros, J. J. (1994) *Mol. Gen. Genet.* **242**, 490–494.
- Oldenburg, K. R., Vo, K. T., Michaelis, S. & Paddon, C. (1997) *Nucleic Acids Res.* **25**, 451–452.
- Raymond, C. K., Pownder, T. A. & Sexson, S. L. (1999) *BioTechniques* **26**, 134–141.
- Ninomiya, Y., Suzuki, K., Ishii, C. & Inoue, H. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 12248–12253.
- Bistis, G. N., Perkins, D. D. & Read, N. D. (2003) *Fungal Genet. Newsl.* **50**, 17–19.
- Borkovich, K. A., Alex, L. A., Yarden, O., Freitag, M., Turner, G. E., Read, N. D., Seiler, S., Bell-Pedersen, D., Paietta, J., Plesofsky, N., et al. (2004) *Microbiol. Mol. Biol. Rev.* **68**, 1–108.
- Galagan, J. E., Calvo, S. E., Borkovich, K. A., Selker, E. U., Read, N. D., Jaffe, D., FitzHugh, W., Ma, L. J., Smirnov, S., Purcell, S., et al. (2003) *Nature* **422**, 859–868.
- Zhou, L. & Marzluf, G. A. (1999) *Biochemistry* **38**, 4335–4341.
- Feng, B. & Marzluf, V. (1998) *Mol. Cell. Biol.* **18**, 3983–3990.
- Yuan, G. H., Fu, Y. H. & Marzluf, G. A. (1991) *Mol. Cell. Biol.* **11**, 5735–5745.
- Kang, S. & Metzberg, R. L. (1990) *Mol. Cell. Biol.* **10**, 5839–5848.
- Fu, Y. H., Kneesi, J. Y. & Marzluf, G. A. (1989) *J. Bacteriol.* **171**, 4067–4070.
- Baum, J. A., Geever, R. & Giles, N. H. (1987) *Mol. Cell. Biol.* **7**, 1256–1266.
- Liu, T. D. & Marzluf, G. A. (2004) *Curr. Genet.* **46**, 213–227.
- Barthelmess, I. B. (1982) *Genet. Res.* **39**, 169–185.
- Bibbins, M., Crepin, V. F., Cummings, N. J., Mizote, T., Baker, K., Mellits, K. H. & Connerton, I. F. (2002) *Mol. Genet. Genom.* **267**, 498–505.
- Bailey, L. A. & Ebbole, D. J. (1998) *Genetics* **148**, 1813–1820.
- Yamashiro, C. T., Ebbole, D. J., Lee, B.-K., Brown, R. E., Bourland, C., Madi, L. & Yanofsky, C. (1996) *Mol. Cell. Biol.* **16**, 6218–6228.
- Linden, H., Ballario, P., Arpaia, G. & Macino, G. (1999) *Adv. Genet.* **41**, 35–54.
- Dunlap, J. C. (1999) *Cell* **96**, 271–290.
- Li, D., Bobrowicz, P., Wilkinson, H. H. & Ebbole, D. J. (2005) *Genetics* **170**, 1091–1104.
- Feng, B., Haas, H. & Marzluf, G. A. (2000) *Biochemistry* **39**, 11065–11073.
- Aramayo, R., Peleg, Y., Addison, R. & Metzberg, R. (1996) *Genetics* **144**, 991–1003.
- Tomita, H., Soshi, T. & Inoue, H. (1993) *Mol. Gen. Genet.* **238**, 225–233.
- Ferreira, A. V. B., An, Z. Q., Metzberg, R. L. & Glass, N. L. (1998) *Genetics* **148**, 1069–1079.
- Staben, C. & Yanofsky, C. (1990) *Proc. Natl. Acad. Sci. USA* **87**, 4917–4921.
- Gritz, L. & Davies, J. (1983) *Gene* **25**, 179–188.
- Staben, C., Jensen, B., Singer, M., Pollock, J., Schechtman, M., Kinsey, J. & Selker, E. (1989) *Fungal Genet. Newsl.* **36**, 79–81.
- Giaever, G., Chu, A. M., Ni, Connelly, C., Riles, L., Veronneau, S., Dow, S., Lucau-Danila, A., Anderson, K., Andre, B., et al. (2002) *Nature* **418**, 387–391.
- Shoemaker, D. D., Lashkari, D. A., Morris, D., Mittmann, M. & Davis, R. W. (1996) *Nat. Genet.* **14**, 450–456.
- Winzeler, E. A., Shoemaker, D. D., Astromoff, A., Liang, H., Anderson, K., Andre, B., Bangham, R., Benito, R., Boeke, J. D., Bussey, H., et al. (1999) *Science* **285**, 901–906.
- Catlett, N. L., Lee, B.-N., Yoder, O. C. & Turgeon, B. G. (2002) *Fungal Genet. Newsl.* **50**, 9–11.
- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997) *Nucleic Acids Res.* **25**, 3389–3402.
- Springer, M. L. (1993) *BioEssays* **15**, 365–374.
- Matsuyama, S. S., Nelson, R. E. & Siegel, R. W. (1974) *Dev. Biol.* **41**, 278–287.
- Bailey-Shrode, L. & Ebbole, D. J. (2004) *Genetics* **166**, 1741–1749.
- Raju, N. B. & Leslie, J. F. (1992) *Genome* **35**, 815–826.
- Masloff, S., Poggeler, S. & Kuck, U. (1999) *Genetics* **152**, 191–199.
- Vienken, K., Scherer, M. & Fischer, R. (2005) *Genetics* **169**, 619–630.
- Chua, G., Robinson, M. D., Morris, Q. & Hughes, T. R. (2004) *Curr. Opin. Microbiol.* **7**, 638–646.
- Paluh, J. L. & Yanofsky, C. (1991) *Mol. Cell. Biol.* **11**.
- Ebbole, D. & Sachs, M. S. (1990) *Fungal Genet. Newsl.* 17–18.
- Arnaise, S., Zickler, D., Poisier, C. & Debuchy, R. (2001) *Mol. Microbiol.* **39**, 54–64.
- Abate-Shen, C. (2002) *Nat. Rev. Cancer* **2**, 777–785.
- Ford, H. L. (1998) *Cell Biol. Int.* **22**, 397–400.
- Mahaffey, J. W. (2005) *Curr. Opin. Genet. Dev.* **15**, 422–429.
- Firulli, A. B. & Thattaliyath, B. D. (2002) in *International Review of Cytology, A Survey of Cell Biology*, Vol. 214, pp. 1–62.
- Martinez, E. (2002) *Plant Mol. Biol.* **50**, 925–947.
- Pickett, F. B. & Meekswagner, D. R. (1995) *Plant Cell* **7**, 1347–1356.
- Krakauer, D. C. & Nowak, M. A. (1999) *Semin. Cell Dev. Biol.* **10**, 555–559.
- Winston, F., Dollard, C. & Ricupero-Hovasse, S. L. (1995) *Yeast* **11**, 53–55.
- Christianson, T. W., Sikorski, R. S., Dante, M., Shero, J. H. & Hieter, P. (1992) *Gene* **110**, 119–122.
- Gera, J. F., Hazbun, T. R. & Fields, S. (2002) *Methods Enzymol.* **350**, 499–512.
- Pall, M. L. (1993) *Fungal Genet. Newsl.* **40**, 58.
- Pall, M. L. & Brunelli, J. P. (1993) *Fungal Genet. Newsl.* **40**, 59–62.
- Case, M. E., Schweizer, M., Kushner, S. R. & Giles, N. H. (1979) *Proc. Natl. Acad. Sci. USA* **76**, 5259–5263.
- Yang, Q., Poole, S. I. & Borkovich, K. A. (2002) *Eukaryot. Cell* **1**, 378–390.
- Davis, R. H. & deSerres, F. J. (1970) *Methods Enzymol.* **71A**, 79–143.
- Ivey, F. D., Hodge, P. N., Turner, G. E. & Borkovich, K. A. (1996) *Mol. Biol. Cell* **7**, 1283–1297.

Corrections

CHEMISTRY. For the article “Tetrahedral structure or chains for liquid water,” by Teresa Head-Gordon and Margaret E. Johnson, which appeared in issue 21, May 23, 2006, of *Proc Natl Acad Sci USA* (103:7973–7977; first published May 12, 2006; 10.1073/pnas.0510593103), the authors note that there was a plotting error in Fig. 3a. The corrected figure and its legend are shown below. This error does not affect the conclusions of the article.




Fig. 3. Comparison of $H_{00}(Q)$ structure factors from experiments reported in ref. 12 (black) against the asymmetric water model (ref. 14; red) (a) and the classical polarizable TIP4P-pol2 model (ref. 16; red) (b). The simulated structure factors using the asymmetric and TIP4P-pol2 models show good agreement with x-ray scattering for $Q > 6.5 \text{ \AA}^{-1}$. However, the simulated structure factors for the asymmetric model show significant disagreement for $Q < 6.5 \text{ \AA}^{-1}$, whereas the TIP4P-pol2 model shows excellent agreement over the full Q range of the measured x-ray data.

www.pnas.org/cgi/doi/10.1073/pnas.0608020103

APPLIED BIOLOGICAL SCIENCES. For the article “Adeno-associated virus serotype 9 vectors transduce murine alveolar and nasal epithelia and can be readministered,” by Maria P. Limberis and James M. Wilson, which appeared in issue 35, August 29, 2006, of *Proc Natl Acad Sci USA* (103:12993–12998; first published August 22, 2006; 10.1073/pnas.0601433103), the authors note that on page 12995, right column, second full paragraph, the third sentence is incorrect in part. “For each vector the highest amount of vector in terms of vector per diploid genome was in the lung [approximately six and one vectors per diploid genome for AAV2/5 and AAV2/9, respectively (Table 1); note that 1.5×10^5 vector genomes per 100 ng of cellular DNA is equivalent to one vector genome per diploid genome of the cell]” should read: “For each vector, the highest amount of vector in terms of vector per diploid genome was in the lung [approximately six and one vectors per diploid genome for AAV2/5 and AAV2/9, respectively (Table 1); note that 1.5×10^4 vector genomes per 100 ng of cellular DNA is equivalent to one vector genome per diploid genome of the cell].” This error does not affect the conclusions of the article.

www.pnas.org/cgi/doi/10.1073/pnas.0608505103

GENETICS. For the article “A high-throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription factors,” by Hildur V. Colot, Gyungsoon Park, Gloria E. Turner, Carol Ringelberg, Christopher M. Crew, Liubov Litvinkova, Richard L. Weiss, Katherine A. Borkovich, and Jay C. Dunlap, which appeared in issue 27, July 5, 2006, of *Proc Natl Acad Sci USA* (103:10352–10357; first published June 26, 2006; 10.1073/pnas.0601456103), the authors note that on page 10354, in lines 13 and 14 of the last paragraph, right column, the gene *mel-1* should be *mld-1*. These errors do not affect the conclusions of the article.

www.pnas.org/cgi/doi/10.1073/pnas.0608497103

MEDICAL SCIENCES. For the article “Bim and Bad mediate imatinib-induced killing of Bcr/Abl⁺ leukemic cells, and resistance due to their loss is overcome by a BH3 mimetic,” by Junya Kuroda, Hamsa Puthalakath, Mark S. Cragg, Priscilla N. Kelly, Philippe Bouillet, David C. S. Huang, Shinya Kimura, Oliver G. Ottmann, Brian J. Druker, Andreas Villunger, Andrew W. Roberts, and Andreas Strasser, which appeared in issue 40, October 3, 2006, of *Proc Natl Acad Sci USA* (103:14907–14912; first published September 22, 2006; 10.1073/pnas.0606176103), the authors note that the affiliation information for Andreas Villunger was incorrect in part. The correct institution name is “Innsbruck Medical University.”

www.pnas.org/cgi/doi/10.1073/pnas.0608594103

